

**Linear vs. Piecewise-Linear
Embeddability of Simplicial
Complexes**

**U. Brehm
K. S. Sarkaria**

**MAX-PLANCK-INSTITUT
FÜR MATHEMATIK
BONN**

Linear vs. Piecewise-Linear Embeddability of Simplicial Complexes

**U. Brehm
K. S. Sarkaria**

U. Brehm
Mathematisches Institut
Technische Universität
Berlin

Germany

Max-Planck-Institut für Mathematik
Gottfried-Claren-Straße 26
D-5300 Bonn 3

Germany

K. S. Sarkaria
Department of Mathematics
Panjab University
Chandigarh 160014

India

MPI / 92-52

§1. Introduction

(1.1) Definitions.

In order to state our results we will first fix the definitions of the notions mentioned in the title.

SIMPLICIAL COMPLEX K : by this we mean a finite set whose members, called its *simplices*, are themselves finite sets, and which is closed under subsets. The members of the simplices of K are called K 's *vertices*.

Its realization K : If K has N vertices, then by thinking of these as the canonical basis vectors of \mathbb{R}^N , and of each simplex as the convex hull of its vertices, one obtains a subspace of \mathbb{R}^N , which too will be denoted K .

LINEAR EMBEDDABILITY OF K IN \mathbb{R}^m : a one-one map $e: K \rightarrow \mathbb{R}^m$ (from this realization K) will be called a *linear embedding* if it is the restriction of a linear map $\mathbb{R}^N \rightarrow \mathbb{R}^m$.

Note that for $m \geq 2(\dim K) + 1$, any general position linear map $\mathbb{R}^N \rightarrow \mathbb{R}^m$ will restrict to such a linear embedding of K in \mathbb{R}^m . Thus the cases of interest are $\dim K \leq m \leq 2(\dim K)$.

PIECEWISE-LINEAR EMBEDDABILITY OF K IN \mathbb{R}^m : this means that, for some $r \geq 0$, the r th derived $K^{(r)}$ of K embeds linearly in \mathbb{R}^m .

Here the r th *derived* is defined inductively by $K^{(0)} = K$ and $K^{(r)} = (K^{(r-1)})'$, where L' denotes the simplicial complex whose simplices are sets of nonempty simplices of L which are totally ordered under \subset .

By mapping each vertex of K' (a simplex of K) to its barycentre, one gets the linear *barycentric* embedding of K' onto K , and so, by iteration, $K^{(r)} \xrightarrow{\cong} K$.

Composing with the inverse of this barycentric subdivision map, each linear embedding $K^{(r)} \rightarrow \mathbb{R}^m$ determines a one-one *piecewise-linear embedding* $e: K \rightarrow \mathbb{R}^m$.

The notion of piecewise-linear embeddability has been much studied – see e.g. Hudson [7] and Rourke-Sanderson [8] which will be our references for all other piecewise-linear terminology – because it avoids the possible wildness of topological embeddings, but is at the same time flexible enough to make it much easier to handle than linear (or ‘simplex-wise-linear’ or ‘geometric’) embeddability.

(1.2) Statements of results.

As an easy consequence of a theorem of Steinitz [14], 1922, it follows that a one-dimensional complex, i.e. a *graph* K^1 , will embed piecewise-linearly (or even topologically) in \mathbb{R}^2 , only if it occurs as a subcomplex of the boundary of a simplicial 3-polytope: so *à fortiori* such a K^1 must also embed linearly in \mathbb{R}^2 . See also Wagner [17], Fáry [3], Stein [13] and Stojaković [15].

In 1969, Grünbaum [6, p.502] conjectured that, likewise, for all $n \geq 2$, the piecewise-linear embeddability of a K^n in \mathbb{R}^{2n} will be sufficient to guarantee its linear embeddability in \mathbb{R}^{2n} . We show that this conjecture is false in the following very strong sense.

Theorem A. *For each $n \geq 2$, $r \geq 0$, there is a simplicial n -complex L which embeds piecewise-linearly in \mathbb{R}^{2n} , but whose r th derived $L^{(r)}$ does not embed linearly in \mathbb{R}^{2n} .*

By virtue of a theorem of van Kampen [16, p.152], 1932, it is known that if K^n is a *pseudomanifold*, i.e. if each of its $(n-1)$ -simplices is incident to at most two n -simplices, then it embeds piecewise-linearly in \mathbb{R}^{2n} . Though the K^n 's of Theorem A are *not* pseudomanifolds, we do have, for ambient dimension one less, the following result which exhibits a similar phenomenon on the part of some ‘higher-dimensional Möbius strips’.

Theorem B. *For each $n = 2^k, k \geq 1$, there is a K^n homeomorphic to M^n , the piecewise-linear manifold-with-boundary obtained by deleting an n -ball B^n from real projective space $\mathbb{R}P^n$, such that K^n embeds piecewise-linearly, but not linearly, in \mathbb{R}^{2n-1} .*

The case $n = 2$ of Theorem B, viz. that of the ordinary Möbius strip, was dealt with by the first author in [2].

Method of proof. The constructions given below to establish Theorems B and A are based on the notion of *linking*, and follow the basic strategy already used in [2].

First, we arrange that, under any arbitrary piecewise-linear embedding, some two spherical subcomplexes will link each other with linking number ≥ 2 .

Second, we take care to triangulate these two spheres by so few vertices that, under a linear embedding, this would be impossible.

We now recall what we need about linking, for more see e.g. Rourke-Sanderson [8], pp. 68-73, and Wu [19], pp. 175-181.

LINKING NUMBER: of any oriented p.l. sphere $S^{a-1} \subset \mathbb{R}^m$, with a disjoint oriented closed p.l. manifold $M^{m-a} \subset \mathbb{R}^m$, is the *intersection number*, i.e. counts the algebraical number of intersections, of any bounding compatibly oriented general position p.l. disk D^a , $\partial D^a = S^{a-1}$, with M^{m-a} . This is done by assigning an orientation to \mathbb{R}^m , and counting each of these intersections as +1 or -1 depending on whether the local orientation of D followed by that of M agrees with that of \mathbb{R}^m or not.

If this number is zero, i.e. if S^{a-1} does not link M^{m-a} , then $S^{a-1} \hookrightarrow M^{m-a}$ extends to a map f of D^a into \mathbb{R}^m such that $f(D^a) \cap M^{m-a} = \emptyset$.

Upto sign, the linking number of $S^{a-1} \subset \mathbb{R}^m$ with a sphere $S^{m-a} \subset \mathbb{R}^m$, is same as that of S^{m-a} with S^{a-1} , and coincides with the *degree* of an associated map - cf. proof of (3.1.1) - of the join $S^m = S^{a-1} \cdot S^{m-a}$ into itself.

§2. Higher Möbius strips

(2.1) Proof of Theorem B.

As is well known the manifold-with-spherical boundary, $M^n = \mathbb{R}P^n - (\text{int}B^n)$, $\partial M^n = \partial B^n = S^{n-1}$, can be considered as a twisted line bundle over a *core* submanifold $\mathbb{R}P^{n-1} \subset M^n$.

(2.1.1) M^n embeds piecewise-linearly in \mathbb{R}^{2n-1} .

To see this we can e.g. first embed (some triangulation of) the core $\mathbb{R}P^{n-1}$ piecewise-linearly in \mathbb{R}^{2n-2} , and so a trivial line bundle over it into \mathbb{R}^{2n-1} . The assertion now follows because we can locally twist the trivial bundle, for each of the \mathbb{R}^{n-1} worth of directions along $\mathbb{R}P^{n-1}$, in the corresponding direction from the \mathbb{R}^{n-1} worth of directions available complementary to the embedded trivial bundle.

(2.1.2) The bounding sphere of M^n links its core under any piecewise-linear embedding $e : M^n \rightarrow \mathbb{R}^{2n-1}$.

We give below, for all $k \geq 2$, a geometric argument; another more algebraical proof is sketched later in (2.2).

Assume, if possible, that $e(S^{n-1})$ does not link $e(\mathbb{R}P^{n-1})$. So we can extend the embedding e to a general position map f (of some triangulation) of $\mathbb{R}P^n$ into \mathbb{R}^{2n-1} , such that $f(\mathbb{R}P^{n-1}) \cap f(B^n) = \emptyset$.

We will now use some well-known constructions – cf. Zeeman [20] and [9] – to modify f to a piecewise-linear embedding g of $\mathbb{R}P^n$ in \mathbb{R}^{2n-1} ; this suffices to furnish the desired contradiction because a theorem of Thom – see e.g. Steenrod [12], p. 34 – tells us that if $n = 2^k$, then $\mathbb{R}P^n$ does not embed in \mathbb{R}^{2n-1} .

We begin by noting that the singularities $\text{sing}(f)$ of f constitute an, at most one-dimensional, subset of the open n -ball $\mathbb{R}P^n - \mathbb{R}P^{n-1}$. So we can find a 2-dimensional conical subset A of this open n -ball such that $A \supset \text{sing}(f)$.

In case $k \geq 3$ one has $3 + n < 2n - 1$, so in this case we can enlarge the 2-dimensional subset $f(A)$ of $f(\mathbb{R}P^n) \subset \mathbb{R}^{2n-1}$ to a 3-dimensional cone $C \subset \mathbb{R}^{2n-1}$ which meets $f(\mathbb{R}P^n)$ only in $f(A)$.

We now choose regular neighbourhoods $N(A)$ of A in $\mathbb{R}P^n$, and $N(C)$ of C in \mathbb{R}^{2n-1} , such that the exterior, boundary, and the interior of $N(A)$ are mapped by f into the exterior, boundary, and the interior, respectively, of $N(C)$. Note that $N(A)$ is an n -ball, while $N(C)$ is a $(2n - 1)$ -ball, and that f is one-one outside $\text{int}(N(A))$. So, by coning $f(\partial(N(A)))$ over an interior point of the ball $N(C)$, we obtain an embedding $g : \mathbb{R}P^n \rightarrow \mathbb{R}^{2n-1}$.

In case $k = 2$ we can, in the first instance, only ensure that the cone C meets $f(\mathbb{R}P^n)$ in finitely many points besides $f(A)$. But then, by using a preliminary modification of f near some one-dimensional tree containing this zero-dimensional singular set, we can replace f by an f' such that C meets $f'(\mathbb{R}P^n)$ only in $f'(A) = f(A)$. After that we proceed as above to modify f' to an embedding g .

(2.1.3) *The image of the bounding sphere of M^n has a nonzero and even self-linking number under any piecewise-linear embedding $e : M^n \rightarrow \mathbb{R}^{2n-1}$.*

Here, by *self-linking number* of $\partial M^n = S^{n-1}$, we mean its linking number with a disjoint isotopic $\Sigma^{n-1} \subset M^n$.

To see the above note that any general position n -disk $D^n \subset \mathbb{R}^{2n-1}$, with $\partial D^n = e(S^{n-1})$, hits the core $e(\mathbb{R}P^{n-1})$ transversely in finitely many points. By (2.1.2) we know that the algebraical number t of such

intersections is nonzero.

Now push S^{n-1} uniformly, along the fibers of the line bundle M^n over $\mathbb{R}P^{n-1}$, to obtain an isotopic sphere Σ^{n-1} arbitrarily close to the core $\mathbb{R}P^{n-1}$. Then the n -disk $D^n \subset \mathbb{R}^{2n-1}$ will intersect this double cover $e(\Sigma^{n-1})$ of $e(\mathbb{R}P^{n-1})$ transversely in $2t$ points.

(2.1.4) CONSTRUCTION OF K^n : Triangulate the boundary S^{n-1} and the isotopic sphere Σ^{n-1} of (2.1.3) as boundaries ∂s^n and $\partial \sigma^n$ of n -simplices s^n and σ^n . We choose any triangulation K^n of M^n which extends – cf. Armstrong [1] – this triangulation $\partial s^n \cup \partial \sigma^n$ of $S^{n-1} \cup \Sigma^{n-1}$. For example one can choose the explicit K^n 's of (2.2.5).

(2.1.5) K^n does not embed linearly in \mathbb{R}^{2n-1} .

Otherwise, there will be some general position linear map $e : \mathbb{R}^N \rightarrow \mathbb{R}^{2n-1}$, whose restriction to the realization K^n is one-one.

The e -images of the closed simplices s^n and σ^n will either not intersect, or intersect in a line segment. In the latter case, if both ends of the line segment lie on the boundary of the same closed simplex, say on $e(\partial(s^n))$, then there is no linking, because $e(s^n) \cap e(\partial \sigma^n) = \emptyset$. And, if the two ends of the line segment lie on different boundaries, then we have $\text{card}(e(s^n) \cap e(\partial \sigma^n)) = 1$.

So the linking number of S^{n-1} and Σ^{n-1} , under a linear embedding e , would be 0 or ± 1 , which contradicts (2.1.3). *q.e.d.*

(2.2) Deleted joins.

Embeddability questions – see e.g. [10] and its references – are intimately related to the following notion.

DELETED JOIN K_* : subcomplex of $K \cdot \overline{K}$, the join of two disjoint copies of K , consisting of all simplices $\sigma \cdot \overline{\theta}$ such that $\sigma \cap \theta = \emptyset$, and equipped with the free \mathbb{Z}_2 -action $\sigma \cdot \overline{\theta} \mapsto \theta \cdot \overline{\sigma}$.

Remarks (2.2.1) - (2.2.3) below sketch an alternative proof of (2.1.2) via deleted joins.

(2.2.1) If $e(S^{n-1})$ were not linking $e(\mathbb{R}P^{n-1})$ under the embedding $e : M^n \rightarrow \mathbb{R}^{2n-1}$, then there would be a continuous \mathbb{Z}_2 -map from the deleted join T_* , of some triangulation of $\mathbb{R}P^n$, into the antipodal $(2n-1)$ -sphere S^{2n-1} .

This is not hard to check, cf. proof of (3.1.3). In fact there would also

be such a \mathbb{Z}_2 -map from the *deleted product* T_* , i.e. the ‘mid-section’ of T_* consisting of all cells $\sigma \times \bar{\theta}$ such that $\sigma \cap \theta = \emptyset$, into the antipodal sphere S^{2n-2} of one dimension less.

(2.2.2) WU LEMMA. *The \mathbb{Z}_2 -homotopy types of the deleted join and the deleted product of a simplicial complex are topological invariants of the space underlying the complex.*

This is harder – cf. Wu [19, Ch.2] for products – but it will be shown in [11] that, with some care, this important fact generalizes even to *higher deleted joins*, i.e. analogues of K_* for groups G other than \mathbb{Z}_2 .

(2.2.3) So, using any convenient triangulation of $\mathbb{R}P^n$, $n = 2^k$, it suffices to show by a calculation of the characteristic classes of the free \mathbb{Z}_2 -homotopy type $(\mathbb{R}P^n)_*$, that there is no continuous \mathbb{Z}_2 -map from it to S^{2n-2} .

This calculation, which will be included in [11], is reminiscent of, but more general than, the proof of the

BORSUK-ULAM THEOREM. *There is no continuous \mathbb{Z}_2 -map from S^p to S^q for $p > q$.*

However for $k = 1$, the Borsuk-Ulam Theorem itself provides the desired contradiction because of the following remarkable fact.

(2.2.4) *The deleted join of the 6-vertex real projective plane $\mathbb{R}P_6^2$ is \mathbb{Z}_2 -homeomorphic to the antipodal 4-sphere.*

We recall that $\mathbb{R}P_6^2$ is a \mathbb{Z}_2 -quotient or, if one prefers, one of the two parts of a yin-yang decomposition — cf. Grothendieck [5] — of the regular 12-vertex 2-sphere, i.e. the ubiquitous *icosahedron*.

The above result is not hard to check. In fact the second author hopes to include in [11] a complete classification of all K^n ’s for which K_* is a closed pseudomanifold. For example, if this pseudomanifold is n -dimensional, then it has to be the *octahedral n -sphere* $(\sigma_n^n)_*$ and — see [10] — if it is $(2n + 1)$ -dimensional, then it has to be a join of some *Flores’ spheres* $(\sigma_{2n-1}^{2n})_*$. Here and below σ_j^i denotes the j -skeleton of an i -simplex.

(2.2.5) *The omission of the n -simplex $\bar{\sigma}^n$, from the simplicial join across σ^n , of any triangulation of $\mathbb{R}P^n$ and the octahedral n -sphere $(\sigma_n^n)_*$, results in a K^n which satisfies the requirements of (2.1.4).*

This is straightforward. Here, by *simplicial join* $\mathbb{R}P^n \# (\sigma_n^n)_*$ across σ^n we mean the operation of first omitting an open n -simplex from the first factor and σ^n from the second factor, and then glueing the remaining complexes together by identifying the boundaries of these n -simplices.

Note in particular that $(\mathbb{R}P_6^2 \# (\sigma_2^2)_*) - \overline{\sigma^2}$ gives the 9-vertex Möbius strip [2] which fails to embed linearly in \mathbb{R}^3 .


(2.2.6) The characteristic class computations of (2.2.3) suggest that if $\alpha(n)$ denotes the number of 1's in the binary expansion of n , then the simplicial Möbius n -strips K^n , $n \geq 2$, of (2.1) embed piecewise-linearly, but not linearly, in the space $\mathbb{R}^{2n-\alpha(n)}$.

§3. Grünbaum's conjecture

(3.1) Proof of Theorem A.

We will first consider the case $n = 2$.

Let $M\ddot{o}_6$ denote the 6-vertex Möbius strip, i.e. $\mathbb{R}P_6^2$ minus one of its 2-simplices which will be called s^2 . We note that, with appropriate orientations, $M\ddot{o}_6$'s boundary ∂s^2 is homologous to twice its core $\partial \sigma^2$, where $\sigma^2 \notin \mathbb{R}P_6^2$ denotes the complementary 2-simplex $vert(\mathbb{R}P_6^2) - s^2$.



Besides $M\ddot{o}_6$, we will also use a disjoint 6-simplex τ^6 , one of whose 2-faces will also be called s^2 , with the complementary 3-simplex $vert(\tau^6) - s^2$ denoted by φ^3 .

(3.1.1) THE 2-COMPLEXES L_t . Each of these will contain a triangle called ∂s^2 . For $t = 0$ we set

$$L_0 = \tau_2^6 - s^2,$$

and having defined L_t , $t \geq 0$, obtain L_{t+1} from L_t by identifying its ∂s^2 with the core $\partial \sigma^2$ of a disjoint copy of $M\ddot{o}_6$. So, after this identification, the boundary ∂s^2 of $M\ddot{o}_6$ becomes the ∂s^2 of L_{t+1} .

(3.1.2) *The 2-complexes L_t embed piecewise-linearly in \mathbb{R}^4 .*

This is clear for $t = 0$.

So, assume inductively that there is a piecewise-linear embedding $e : L_t \rightarrow \mathbb{R}^4$, for some $t \geq 0$. Since $M\ddot{o}_6$ embeds piecewise-linearly even in \mathbb{R}^3 , we can extend e to a general position piecewise-linear map $f : L_{t+1} \rightarrow \mathbb{R}^4$, with its finitely many double points (x, y) all such that $x \in L_t$ and $y \in M\ddot{o}_6$. For each such y choose a disjoint arc of $M\ddot{o}_6$ from y to its boundary ∂s^2 . Removing from L_{t+1} small regular neighbourhoods of all these arcs we get a subspace X piecewise-linearly homeomorphic to L_{t+1} on which the map f is one-one.

(3.1.3) *The disjoint spheres $\partial\varphi^3$ and ∂s^2 of L_0 must link under any piecewise-linear embedding $e : L_0 \rightarrow \mathbb{R}^4$.*

By a lemma of Flores [4] the deleted join $(\tau_2^6)_*$ is an antipodal 5-sphere. So Borsuk-Ulam tells us that there can not be a continuous \mathbb{Z}_2 -map from it to S^4 .

But, S^4 has the same \mathbb{Z}_2 -homotopy type as the join $\mathbb{R}^4 \cdot \overline{\mathbb{R}^4}$ minus its diagonal, i.e. all points of the type $\frac{1}{2}x + \frac{1}{2}\bar{x}$. And, there is a continuous \mathbb{Z}_2 -map of $(L_0)_*$ into this space, viz. the map e_* defined by

$$\lambda x + (1 - \lambda)\bar{y} \mapsto \lambda e(x) + (1 - \lambda)\overline{e(y)}.$$

The closure of $(\tau_2^6)_* - (L_0)_*$ consists of the 5-ball $\partial\varphi^3 \cdot \overline{s_2^2}$ and its conjugate. The restriction of e_* to the boundary of this 5-ball has degree zero iff the linking number of the spheres $e(\partial\varphi^3)$ and $e(\partial s^2)$ is zero. So, if this were the case, e_* would extend to yield a continuous \mathbb{Z}_2 -map $(\tau_2^6)_* \rightarrow S^4$, which is not possible.

(3.1.4) *The disjoint spheres $\partial\varphi^3$ and ∂s^2 of L_t , $t \geq 0$, must have linking number at least 2^t (in absolute value) under any piecewise-linear embedding $e : L_t \rightarrow \mathbb{R}^4$.*

We argue by induction starting from the above case $t = 0$. The triangle ∂s^2 of complex L_t , $t \geq 1$, is homologous to twice the triangle $\partial\sigma^2 \subset M\ddot{o}_6$ which was identified (3.1.1) to the triangle ∂s^2 of L_{t-1} to form L_t . So each transverse intersection under e of the latter, with a general position 3-disk spanning $e(\partial\varphi^3)$, gives rise to two intersections of the former having the same intersection number.

(3.1.5) For any $r \geq 0$ we can choose t so big that the r th derived of $L = L_t$ does not embed linearly in \mathbb{R}^4 .

The number of simplices, contained in the simplicial 2 and 1-spheres occuring as the r th deriveds of $\partial\varphi^3$ and ∂s^2 , is bounded in terms of r . From this it follows easily that, under any linear embedding of the union of these spheres in \mathbb{R}^4 , the absolute value of the linking number is also bounded by a constant depending only on r . Choose any t such that 2^t is bigger than this number and use (3.1.4).

This concludes the proof of Theorem A for $n = 2$.

(3.1.6) For $n \geq 3$ the above argument modifies as follows :

(a) Instead of $M\ddot{o}_6$ we use its $(n-3)$ -fold suspension $S^{n-3}(M\ddot{o}_6)$. Note that in it the $(n-1)$ -sphere $S^{n-3}(\partial s^2)$ is homologous to twice the $(n-1)$ -sphere $S^{n-3}(\partial\sigma^2)$.

(b) The n -complexes $L_{n,t}$, $t \geq 0$, are defined almost as before except for one small change. Instead of the n -skeleton of a τ^{2n+2} , minus one n -face u^n , we start with

$$L_{n,0} = (\tau_n^{2n+2} - u^n) \cup A^n,$$

where A^n is a simplicial annulus $S^{n-1} \times I$ having boundary $\partial A^n = \partial u^n \cup S^{n-3}(\partial s^2)$. So we have a $S^{n-3}(\partial s^2)$ in $L_{n,0}$ which is homologous to ∂u^n . For any $t \geq 1$, we now obtain $L_{n,t}$ from $L_{n,t-1}$ by identifying this $S^{n-3}(\partial s^2)$ of $L_{n,t-1}$, with the $S^{n-3}(\partial\sigma^2)$ of a disjoint copy of $S^{n-3}(M\ddot{o}_6)$.

The rest of the argument is unchanged: the piecewise-linear embeddability of these n -complexes in \mathbb{R}^{2n} follows just as in (3.1.2), and the same argument as in (3.1.3) shows that the disjoint spheres $\partial\varphi^{n+1}$ and ∂u^n of $L_{n,0}$ link under any embedding in \mathbb{R}^{2n} , from which it follows almost as before that the linking number of $\partial\varphi^{n+1}$ and $S^{n-3}(\partial s^2)$ is $\geq 2^t$ for any embedding of $L_{n,t}$ in \mathbb{R}^{2n} ... q.e.d.

(3.2) Concluding remarks.

We will now consider some variations of the above construction which give in particular a generalization (3.2.3) of Theorem A and a corollary (3.2.5) pertaining to linear immersions.

(3.2.1) Examples $L_{n,t}$ analogous to those of (3.1) can be made starting from any Kuratowski n -complex [9]

$$T^n = \tau_{n_1-1}^{2n_1} \cdot \tau_{n_2-1}^{2n_2} \cdot \dots \cdot \tau_{n_k-1}^{2n_k}, \quad n_1 + \dots + n_k = n + 1,$$

instead of just τ_n^{2n+2} .

For instance had we started off by setting $L_0 = \tau_1^4 \cdot \tau_0^2 - s^1 \cdot s^0$, then the analogue of (3.1.3) is that the 2-sphere $\partial\phi^2 \cdot \partial\phi^1$, formed by the vertices of L_0 not in the omitted 2-simplex $s^1 \cdot s^0$, always links the boundary of $s^1 \cdot s^0$ under any embedding of L_0 into \mathbb{R}^4 .

(3.2.2) *Analogous constructions also give some n -complexes $L_{n,m,t}$ which embed piecewise-linearly, but not linearly in \mathbb{R}^m , for some other n 's and m 's such that $n < m < 2n$.*

We now start with different T^n 's. For example, we can start with the join of $m-n$ disjoint copies of τ_0^2 (i.e. three points) and $2n-m$ disjoint copies of τ_0^0 (i.e. one point). Then the deleted join T_* is an antipodal $(m+1)$ -sphere; so there is no \mathbb{Z}_2 -map from it to S^m . Omitting an n -face from this T^n and proceeding as in (3.1.6) gives such complexes.

Their piecewise-linear embeddability in \mathbb{R}^m follows from arguments analogous to those of (3.1.2) which remain valid at least under conditions like $m \geq \frac{3}{2}n + 1$ - cf. [18] - and thus we obtain examples of the above sort.

(3.2.3) *For each $n \geq 2$, $r \geq 1$, $n < m \leq 2n$, there is a simplicial n -complex which embeds piecewise-linearly in \mathbb{R}^m , but whose r th derived does not embed linearly in \mathbb{R}^m .*

Furthermore, if $n \geq 3$, we can take $n \leq m \leq 2n$ in the above.

These generalizations of Theorem A follow by using (3.2.2): e.g. one takes disjoint union of an $L_{[\frac{m}{2}],m,t}$ and a σ_n^n , etc.

We note that a finesse is required when dealing with the case $n = 2$, $m = 3$ of (3.2.3) since, by attaching Möb's à la (3.1.1), one now loses piecewise-linear embeddability. To overcome this, attach instead, at each step, an $\mathbb{R}P_6^2$ minus a 2-simplex s^2 having exactly one vertex on the attaching triangle $\partial\sigma^2$.

(3.2.4) *By iterating the construction (3.1.1) indefinitely one obtains an infinite 2-complex L_∞ , which embeds topologically, but not piecewise-linearly, in \mathbb{R}^4 .*

This is clear. Here, by *topologically embeddable*, we mean simply that there exists a continuous one-one map from L_∞ into \mathbb{R}^4 .

Construction of such *finite* complexes is much harder, but might be implicit in the well-known work of R.D.Edwards and M.H.Freedman.

(3.2.5) For each $n \geq 3$, $r \geq 0$, $\max\{n, 4\} \leq m < 2n$, there is a simplicial n -complex which embeds piecewise-linearly in \mathbb{R}^m , but whose r th derived does not even immerse linearly in \mathbb{R}^m .

This follows either by considering cones over suitable examples from (3.2.3) or formulating an analogue of (3.2.3) for embeddings in S^m .

(3.2.6) *Embeddability of \mathcal{K} in \mathbb{R}^m .* Thinking again, as in §1, of the N vertices of K , as the canonical basis vectors of \mathbb{R}^N , one gets a bigger (non-compact) space \mathcal{K} , if with each simplex of K is associated the *affine hull* of its vertices in \mathbb{R}^N instead of the convex hull of its vertices.

Note that \mathcal{K} collapses to K , from which it follows that the topological embeddability of K in \mathbb{R}^m implies that of \mathcal{K} . But it is very easy to see – e.g. consider a segment and a disjoint point in \mathbb{R}^1 – that the *linear* embeddability of \mathcal{K} in \mathbb{R}^m is a strictly stronger notion than that of K .

There will be included in Chapter IV (on “Linear Embeddability”) of [11] some interesting results involving this stronger notion, which incidentally makes sense not only for an ordered field like \mathbb{R} , but for any field whatsoever.

ACKNOWLEDGEMENTS

We would like to thank G.Schild for a question to the first author which led to the present stronger formulation of Theorem A in terms of the non-linear embeddability of a given derived of L , rather than of just L itself.

The second author would like to thank the Max-Planck-Institut für Mathematik, Bonn, and the Technische Universität, Berlin, for making this collaboration possible.

REFERENCES

- [1] M.A.ARMSTRONG, Extending triangulations, *Proc. Am. Math. Soc.* 18 (1967), 701-704.
- [2] U.BREHM, A nonpolyhedral triangulated Möbius strip, *Proc. Am. Math. Soc.* 89 (1983), pp.519-522.
- [3] I.FÁRY, On straight line representations of planar graphs, *Acta. Sci. Math. Szeged* 11 (1948), 229-233.
- [4] A.FLORES, Über n -dimensionale Komplexe die im R_{2n+1} absolut selbstverschlungen sind, *Ergeb. Math. Kolloq.* 6 (1933/34), pp. 4-7.

- [5] A.GROTHENDIECK, "Les Portes sur l'Univers", in, *Récoltes et Semailles*, pp. PU 116-122, Université de Montpellier (1985).
- [6] B.GRÜNBAUM, Imbeddings of simplicial complexes, *Comm. Math. Helv.* 45 (1970), 502-513.
- [7] J.F.P.HUDSON, *Piecewise Linear Topology*, Benjamin, New York (1969).
- [8] C.P.ROURKE and B.J.SANDERSON, *Introduction to Piecewise Linear Topology*, Springer, Berlin (1972).
- [9] K.S.SARKARIA, Embedding and unknotting of some polyhedra, *Proc. Am. Math. Soc.* 100 (1987), pp. 201-203.
- [10] —, Kuratowski complexes, *Topology* 30 (1991), pp. 67-76.
- [11] —, *Van Kampen Obstructions*, book under preparation.
- [12] N.E.STEENROD (notes by D.B.A.EPSTEIN), *Cohomology Operations*, Annals Studies no. 50, Princeton (1962).
- [13] S.K.STEIN, *Convex maps*, *Proc. Amer. Math. Soc.* 2 (1951), 464-466.
- [14] E.STEINITZ, "Polyeder und Raumeinteilungen", in, *Enzykl. math. Wiss.*, vol. 3, part 3A12 (1922), 1-139.
- [15] M.STOJAKOVIĆ, Über die Konstruktion der ebenen Graphen, *Univ. Beograd. Godisnjak Filozof. Fak. Novom Sadu* 4 (1959), 375-378.
- [16] E.R.VAN KAMPEN, Komplexe in euklidischen Räumen, *Abh. Math. Sem.* 9 (1932), pp. 72-78, 152-153.
- [17] K.WAGNER, Bemerkungen zum Vierfarbenproblem, *Jber. Deut. Math.-Verein* 46 (1936), 26-32.
- [18] C.WEBER, Plongements de polyèdres dans le domaine métastable, *Comm. Math. Helv.* 42 (1967), pp. 1-27.
- [19] W.-T.WU, *A theory of imbedding, immersion, and isotopy of polytopes in a Euclidean space*, Science Press, Peking (1965).
- [20] E.C.ZEEMAN, "Polyhedral n -manifolds: II. Embeddings", in, *Topology of 3-Manifolds and Related Topics* (M.K.Fort ed.), Prentice-Hall, N.J. (1961), pp. 64-70.

U.Brehm, Mathematisches Institut, Technische Universität, Berlin, GERMANY.

K.S.Sarkaria, Department of Mathematics, Panjab University, Chandigarh 160014, INDIA.

Current Address: Max-Planck-Institut für Mathematik, Gottfried-Claren-Strasse 26, 5300 Bonn 3, GERMANY.

